

Extended Breastfeeding Fact Sheet

Nursing toddlers benefit NUTRITIONALLY

Although there has been little research done on children who breastfeed beyond the age of two, the available information indicates that breastfeeding continues to be a valuable source of nutrition and disease protection for as long as breastfeeding continues.

"Breast milk continues to provide substantial amounts of key nutrients well beyond the first year of life, especially protein, fat, and most vitamins." (Dewey 2001)

In the second year (12-23 months), 448 mL of breastmilk provides: 29% of energy requirements, 43% of protein requirements, 36% of calcium requirements, 75% of vitamin A requirements, 76% of folate requirements, 94% of vitamin B₁₂ requirements & 60% of vitamin C requirements (Dewey 2001).

Studies done in rural Bangladesh have shown that breastmilk continues to be an important source of vitamin A in the second and third year of life (Persson 1998).

It's not uncommon for weaning to be recommended for toddlers who are eating few solids. However, this recommendation is *not* supported by research. According to Sally Kneidel in "Nursing Beyond One Year" (Kneidel 1990):

"Some doctors may feel that nursing will interfere with a child's appetite for other foods. Yet there has been no documentation that nursing children are more likely than weaned children to refuse supplementary foods. In fact, most researchers in Third World countries, where a malnourished toddler's appetite may be of critical importance, recommend continued nursing for even the severely malnourished (Briend et al, 1988; Tangermann, 1988; Shattock and Stephens, 1975; Whitehead, 1985). Most suggest helping the malnourished older nursing child not by weaning but by supplementing the mother's diet to improve the nutritional quality of her milk (Ahn and MacLean, 1980; Jelliffe and Jelliffe, 1978) and by offering the child more varied and more palatable foods to improve his or her appetite (Tangermann, 1988; Underwood, 1985)."

Nursing toddlers are SICK LESS OFTEN

The American Academy of Family Physicians notes that children weaned before two years of age are at increased risk of illness (AAFP 2001).

Nursing toddlers between the ages of 16 and 30 months have been found to have fewer illnesses and illnesses of shorter duration than their non-nursing peers (Gulick 1986).

"Antibodies are abundant in human milk throughout lactation" (Hamosh 1991; p. 134). In fact, some of the immune factors in breastmilk increase in concentration during the second year and also during the weaning process (Goldman 1983, Goldman & Goldblum 1983, Hamosh 1991).

Per the World Health Organization, *"a modest increase in breastfeeding rates could prevent up to 10% of all deaths of children under five: Breastfeeding plays an essential and sometimes underestimated role in the treatment and prevention of childhood illness."*

Nursing toddlers are SMART

Extensive research on the relationship between cognitive achievement (IQ scores, grades in school) and breastfeeding has shown the greatest gains for those children breastfed the longest.

Nursing toddlers have FEWER ALLERGIES

Many studies have shown that one of the best ways to prevent allergies and asthma is to breastfeed exclusively for at least 6 months and to continue breastfeeding long-term after that point. Breastfeeding can be helpful in preventing allergy by:

- (1) reducing exposure to potential allergens (when baby is exposed later, allergic reactions are less likely),
- (2) speeding maturation of the protective barrier in baby's intestines,
- (3) coating the gut and providing a barrier to potentially allergenic molecules,
- (4) providing anti-inflammatory properties that reduce the risk of infections (which can act as allergy triggers).

Nursing toddlers are WELL ADJUSTED SOCIALLY

According to Sally Kneidel in "Nursing Beyond One Year" (Kneidel 1990): *"Research reports on the psychological aspects of nursing are scarce. One study that dealt specifically with babies nursed longer than a year showed a significant link between the duration of nursing and mothers' and teachers' ratings of social adjustment in six- to eight-year-old children (Ferguson et al, 1987). In the words of the researchers, 'There are statistically significant tendencies for conduct disorder scores to decline with increasing duration of breastfeeding.'"*

Children who achieve independence at their own pace are more secure in that independence than children forced into independence prematurely. Per Elizabeth N. Baldwin, Esq. in "Extended Breastfeeding and the Law": *"Breastfeeding is a warm and loving way to meet the needs of toddlers and young children. It not only perks them up and energizes them; it also soothes the frustrations, bumps and bruises, and daily stresses of early childhood. In addition, nursing past infancy helps little ones make a gradual transition to childhood... Meeting a child's dependency needs is the key to helping that child achieve independence. And children outgrow these needs according to their own unique timetable."*

Nursing a toddler is NORMAL

The American Academy of Pediatrics recommends that *"Breastfeeding should be continued for at least the first year of life and beyond for as long as mutually desired by mother and child... Increased duration of breastfeeding confers significant health and developmental benefits for the child and the mother... There is no upper limit to the duration of breastfeeding and no evidence of psychologic or developmental harm from breastfeeding into the third year of life or longer."* (AAP 2005)

The American Academy of Family Physicians recommends that breastfeeding continue throughout the first year of life and that *"Breastfeeding beyond the first year offers considerable benefits to both mother and child, and should continue as long as mutually desired."* They also note *"If the child is younger than two years of age, the child is at increased risk of illness if weaned."* (AAFP 2001)

A US Surgeon General has stated that it is a lucky baby who continues to nurse until age two. (Novello 1990)

The World Health Organization & UNICEF emphasize the importance of nursing up to two years of age or beyond (WHO 1992, 2002).

Scientific research by Katherine A. Dettwyler, PhD shows that 2.5 to 7.0 years of nursing is what our children have been designed to expect (Dettwyler 1995).

MOTHERS also benefit from nursing past infancy

- Extended nursing delays the return of fertility in some women by suppressing ovulation.
 - Breastfeeding reduces the risk of breast cancer. Studies have found a significant inverse association between duration of lactation and breast cancer risk.
 - Breastfeeding reduces the risk of ovarian cancer.
 - Breastfeeding reduces the risk of uterine cancer.
 - Breastfeeding reduces the risk of endometrial cancer.
- Breastfeeding reduces the risk of rheumatoid arthritis.
 - Breastfeeding protects against osteoporosis. During lactation a mother may experience decreases of bone mineral. A nursing mom's bone mineral density may be reduced in the whole body by 1 to 2 percent while she is still nursing. This is gained back, and bone mineral density may actually increase, when the baby is weaned from the breast. This is not dependent on additional calcium supplementation in the mother's diet.
 - Breastfeeding has been shown to decrease insulin requirements in diabetic women.
 - Breastfeeding mothers tend to lose weight easier.

REFERENCES

Only specifically cited references are included in this list. See www.kellymom.com for additional references.

Ahn CH, MacLean WC Jr. Growth of the exclusively breast-fed infant. *Am J Clin Nutr.* 1980 Feb;33(2):183-92.

American Academy of Family Physicians. AAFP Policy Statement on Breastfeeding. 2001.

American Academy of Pediatrics Section on Breastfeeding. Breastfeeding and the Use of Human Milk. *Pediatrics.* Feb 2005;115(2):496-50.

Baldwin, EN. Extended Breastfeeding and the Law. *Mothering.* 1993 (Spring);66:88.

Briend A, Wojtyniak B and Rowland MG. Breast feeding, nutritional state, and child survival in rural Bangladesh. *Br Med J (Clin Res Ed).* 1988 Mar 26;296(6626):879-82.

Dettwyler KA. "A Time to Wean" from *Breastfeeding: Biocultural Perspectives* (Stuart-Macadam, P. and Dettwyler, K., ed.), New York: Walter de Gruyter, Inc., 1995, p. 305-345.

Dewey KG. Nutrition, Growth, and Complementary Feeding of the Breastfed Infant. *Pediatr Clin North Am.* 2001 Feb;48(1):87-104.

Ferguson DM et al. Breastfeeding and subsequent social adjustment in six- to eight-year-old children. *J Child Psychol Psychiatr Allied Discip.* 1987; 28:378-86.

Goldman AS et al. Immunologic components in human milk during weaning. *Acta Paediatr Scand.* 1983 Jan;72(1):133-4.

Goldman AS, Goldblum RM, Garza C. Immunologic components in human milk during the second year of lactation. *Acta Paediatr Scand.* 1983 May;72(3):461-2.

Gulick EE. The effects of breastfeeding on toddler health. *Pediatr Nurs.* 1986 Jan-Feb;12(1):51-4.

Hamosh M, Dewey, Garza C, et al: *Nutrition During Lactation.* Institute of Medicine, Washington, DC, National Academy Press, 1991.

Jelliffe DB and Jelliffe EF. The volume and composition of human milk in poorly nourished communities. A review. *Am J Clin Nutr.* 1978 Mar;31(3):492-515.

Lawrence R and Lawrence R. *Breastfeeding: A Guide for the Medical Profession*, 5th ed. St. Louis: Mosby, 1999.

Mohrbacher N and Stock J. *The Breastfeeding Answer Book*, Third Revised ed. Schaumburg, Illinois: La Leche League International, 2003.

Kneidel S. Nursing Beyond One Year. *New Beginnings.* 1990 Jul-Aug;6(4):99-103.

Novello A, MD, US Surgeon General, "You Can Eat Healthy," *Parade Magazine* (11 Nov 1990): 5.

Persson V, Greiner T, Islam S, Gebre-Medhin M. The Helen Keller international food-frequency method underestimates vitamin A intake where sustained breastfeeding is common. *Food and Nutrition Bulletin*, vol.19 no.4. Tokyo, Japan: United Nations University Press, 1998.

Riordan J and Auerbach K. *Breastfeeding and Human Lactation*, 2nd ed. Boston and London: Jones and Bartlett, 1999.

Shattock FM, Stephens AJ. Letter: Duration of breast-feeding. *Lancet.* 1975 Jan 11;1(7898):113-4.

Tangermann RH, Rohde JE, et al. Breastfeeding beyond twelve months. *Lancet.* 1988 Oct 29;2(8618):1016.

Underwood BA. Weaning practices in deprived environments: the weaning dilemma. *Pediatrics.* 1985 Jan;75(1 Pt 2):194-8.

Whitehead RG. The human weaning process. *Pediatrics.* 1985 Jan;75(1 Pt 2):189-93.

World Health Organization. Global strategy on infant and young child feeding (Document A55/15). 16 April 2002.

World Health Organization. Innocenti Declaration: World Declaration and Plan of Action for Nutrition. International Conference on Nutrition (ICN), Rome, December 1992. Food and Agricultural Organization and World Health Organization.

World Health Organization. Reducing Mortality from Major Killers of Children: Fact Sheet No 178. World Health Organization Division of Child Health and Development, Geneva, revised 1998.